arXiv:1103.6081v1 [hep-th] 31 Mar 2011

Yukawa Institute for Theoretical Physics
Kyoto University

YITP-11-41

Gregory-Laflamme instability of a slowly rotating black string

Chul-Moon Yoo,[1, *] Sugure Tanzawa,[1, †] and Misao Sasaki[1, 2, ‡]

[1] *Yukawa Institute for Theoretical Physics, Kyoto University Kyoto 606-8502, Japan*
[2] *Korea Institute for Advanced Study 207-43 Cheongnyangni 2-dong,*
Dongdaemun-gu, Seoul 130-722, Republic of Korea

We study the Gregory-Laflamme instability of a 5-dimensional slowly rotating black string in which the 4-dimensional section is described by the Kerr black hole. We treat the rotation in a perturbative way introducing a small parameter for the rotation. It is found that rotation makes the Gregory-Laflamme instability stronger. Both the critical wavelength at the onset of instability and the growth time-scale are found to decrease as the rotation increases.

---

[*] Electronic address: yoo@yukawa.kyoto-u.ac.jp
[†] Electronic address: tanzawa@yukawa.kyoto-u.ac.jp
[‡] Electronic address: misao@yukawa.kyoto-u.ac.jp

I. INTRODUCTION

The Gregory-Laflamme(GL) instability is an s-wave instability for black p-brane solutions [1, 2]. This instability is often quoted in the context of gravitational instabilities of black objects. For instance, the gravitational instability of ultra-rotating Myers-Perry(MP) black holes can be regarded as the one similar to the GL instability [3]. The nonlinear dynamical evolution of the GL instability has been numerically simulated in Ref. [4] but as for the final state it is still under discussion. From view points of the membrane paradigm [5] or field theory/gravity correspondence [6–8], dual or analog fluid models of a black string and the GL instability have been attracting much attention in recent years [9–11].

For the deeper understanding of gravitational instabilities and the duality with fluid models, it is useful to investigate responses to the variation or addition of physical parameters such as the electric charge, the number of dimensions and so on. One of the most fundamental and straightforward extension is to add rotation. In this paper, we consider the $(D+1)$-dim spacetime given by

$$[D\text{−dim MP black hole}] \times \mathbb{R}\,, \tag{1}$$

and in particular focus on the $D = 4$ case. We call these spacetimes *MP black strings* for $D > 4$ and the *Kerr black string* for $D = 4$.

Effects of rotation on the GL instability have been discussed by several authors [3, 10, 12]. The results of these previous studies suggest that, in general, rotation makes the GL instability stronger. We briefly summarize the known facts about the effects of rotation on the GL instability or instabilities on dual/analog fluid models.

- **Ultra-spinning MP black strings [10]**

  When we consider the case of $D > 5$ and only one rotational axis, the angular momentum of the MP black hole is unbounded. For a very large angular momentum, the metric reduces to [13]

  $$[D\text{−dim MP blackhole}] \times \mathbb{R} \sim [(D-2)\text{−dim Schwraschild}] \times \mathbb{R}^3. \tag{2}$$

  Since this is precisely the geometry considered by Gregory and Laflamme [1], it is unstable. Therefore, we can expect that the GL instability persists even if rotation is added.

- **Co-homogeneity-1 MP black strings**

  MP black holes with all the angular momenta being equal in magnitude have the co-homogeneity-1 geometry (ie, the hypersurface of a constant radial coordinate is homogeneous) in odd dimensions. Therefore, when we consider the geometry given by [co-homogeneity-1 MP]×$\mathbb{R}$, it also has the co-homogeneity-1 symmetry in D dimensions as in the case of non-rotating black strings. We call these spacetimes co-homogeneity-1 MP black strings. In Ref. [12], from an analysis of stationary perturbations on co-homogeneity-1 MP black strings, it was predicted that the GL instability persists until extremality. In Ref. [3], the dispersion relation of the instability was numerically calculated and the prediction in

Ref. [12] was confirmed. The threshold wavelength and the instability time-scale become shorter as the rotation increases, that is, rotation makes the GL instability stronger for co-homogeneity-1 MP black strings.

- **Rayleigh-Plateau instability[14]**

  The Rayleigh-Plateau instability in a rotating fluid tube was discussed in Ref. [14] as a system dual to an MP black string. It is reported that the instability strength and threshold wavenumber increase as the rotation increases.

- **Self gravitating cylinder [15]**

  We can find a similar destabilizing effect of rotation on a self-gravitating cylinder in the 4-dimensional Newtonian gravity [15].

In this paper, we consider the Kerr black string with $D = 4$. In this case, since Kerr black hole is not co-homogeneity-1, it is very difficult to solve full perturbation equations. Thus, we treat the rotation in a perturbative way by introducing a small parameter representing the rotation, that is, we consider only slow rotating black strings. We find that the rotation in the present case also makes the GL instability stronger.

This paper is organized as follows. In Sec. II, we give a thermo-dynamical prediction comparing the entropy of a Kerr black string with that of a 5-dim MP black hole. Equations for the metric perturbations to second order in the rotation parameter are derived in Sec. III and the results of the numerical calculation are presented in Sec. IV. Some messy expressions are deferred to Appendices B and C. A minimum review of spherical harmonics is given in Appendix A. We set the $D+1$-dimensional gravitational constant and the speed of light equal to unity.

II. THERMODYNAMICS

First, we discuss the instability of stationary black objects from the thermo-dynamical point of view. We consider a Kerr black string with the length L along the z-direction and compare this with the 5-dimensional MP black hole which has the same total mass and the angular momentum as the Kerr black string. Since the horizon area is regarded as the entropy in black hole thermodynamics, if the horizon area of the Kerr black string is smaller than that of the MP black hole, the Kerr black string is thermodynamically less favorable than the MP black hole. Then, it is expected that the Kerr black string becomes unstable to a perturbation with the wavelength L along the z-direction.

The metric of a Kerr black string is written in the form,

$$\begin{aligned} ds^2 &= -\frac{\triangle - a^2\sin^2\theta}{\Sigma}dt^2 - 2a\sin^2\theta\frac{r^2+a^2-\triangle}{\Sigma}dtd\phi \\ &\quad +\frac{(r^2+a^2)^2-\triangle a^2\sin^2\theta}{\Sigma}\sin^2\theta d\phi^2+\frac{\Sigma}{\triangle}dr^2+\Sigma d\theta^2+dz^2, \end{aligned} \tag{3}$$

where $0 \leq z \leq L$ with $z = 0$ and $z = L$ being identified, and

$$\Sigma = r^2 + a^2\cos^2\theta, \tag{4}$$

$$\triangle = r^2 - 2Mr + a^2 = (r - r_+)(r - r_-). \tag{5}$$

The z =const., 4-dimensional section is given by the metric of a Kerr black hole. The tonal mass E and the total angular momentum J of the Kerr black string are given by

$$E = LM\,, \quad J = Ea\,. \tag{6}$$

The horizon area is given by

$$\frac{A_{\rm BS}}{E^{3/2}} = \frac{4\pi(r_+^2+a^2)L}{E^{3/2}} = 8\pi\frac{1}{l}\left(1+\sqrt{1-j^2l^2}\right), \tag{7}$$

where l and j are non-dimensional quantities defined by

$$l = \frac{L}{E^{1/2}}\,, \quad j = \frac{J}{E^{3/2}}\,. \tag{8}$$

The metric of an MP black hole with single rotation can be written as

$$\begin{aligned} ds^2 &= -dt^2 + \frac{M_{\rm MP}}{\Sigma}\left(dt^2 - a_{\rm MP}\sin^2\theta d\phi\right)^2 + \frac{\Sigma}{\triangle_{\rm MP}}dr^2 \\ &+ \Sigma d\theta^2 + (r^2+a_{\rm MP}^2)\sin^2\theta d\phi^2 + r^2\cos^2\theta dw^2, \end{aligned} \tag{9}$$

where

$$\triangle_{\rm MP} = r^2 + a_{\rm MP}^2 - M_{\rm MP}\,. \tag{10}$$

The total mass and the angular momentum are given by

$$E = \frac{3}{8}\pi M_{\rm MP}\,, \quad J = \frac{2}{3}Ea_{\rm MP}\,. \tag{11}$$

The horizon area is given by

$$\frac{A_{\rm MP}}{E^{3/2}} = \frac{2\pi^2 M_{\rm MP} r_0}{E^{3/2}} = \frac{16}{3}\pi\sqrt{\frac{8}{3\pi} - \frac{9}{4}j^2}\,, \tag{12}$$

where we have assumed $M_{\rm MP} > a_{\rm MP}^2$ and r_0 is the horizon radius given by

$$r_0 = \sqrt{M_{\rm MP} - a_{\rm MP}^2}\,. \tag{13}$$

We define the critical length $L_{\rm crit}$ by the equality of the horizon area, that is,

$$A_{\rm BS} = A_{\rm MP} \ \text{ for } \ L = L_{\rm crit}. \tag{14}$$

Then, fixing the total mass E, we find

$$\begin{cases} A_{\rm BS} \geq A_{\rm MP} \ \text{ for } \ L \leq L_{\rm crit} \\ A_{\rm BS} < A_{\rm MP} \ \text{ for } \ L > L_{\rm crit} \end{cases}, \tag{15}$$

where

$$\frac{L_{\rm crit}}{\sqrt{E}} = \frac{3\sqrt{3\pi}}{16}\sqrt{32 - 27j^2\pi} =: l_{\rm crit}. \tag{16}$$

Hence we expect that Kerr black strings are unstable against a perturbation with the wavelength $L > L_{\rm crit}$. The fact that the value of $l_{\rm crit}$ is a decreasing function of j suggests that the critical wavelength of the GL instability becomes shorter, and hence the critical wavenumber $\mu_{\rm crit}$ becomes larger if we add rotation. In the subsequent sections, by calculating the metric perturbations explicitly, we confirm this observation.

III. PERTURBATION EQUATIONS

To avoid all possible complexities due to the coordinate singularity on the horizon, we use the metric in the Kerr-Schild coordinates given by

$$\begin{aligned} ds^2 &= -\frac{\triangle - a^2\sin^2\theta}{\Sigma}dv^2 + 2dvdr - \frac{2a\sin^2\theta(r^2+a^2-\triangle)}{\Sigma}dvd\chi - 2a\sin^2\theta d\chi dr \\ &\quad + \frac{(r^2+a^2)^2 - \triangle a^2\sin^2\theta}{\Sigma}\sin^2\theta d\chi^2 + \Sigma d\theta^2 + dz^2. \end{aligned} \tag{17}$$

In Einstein gravity, the vacuum field equations are given by

$$R_{\mu\nu} = 0\,. \tag{18}$$

Here, we consider linear metric perturbations on the Kerr black string background. The metric may be decomposed as

$$g = g^{(0)} + \hat{h} + \cdots\,, \tag{19}$$

where $g^{(0)}$ is the metric of the Kerr black string.

Imposing the transverse traceless gauge condition,

$$\nabla_\mu \hat{h}^\mu{}_\nu = 0, \quad \hat{h}^\mu{}_\mu = 0, \tag{20}$$

we can rewrite the metric perturbation equations as

$$\triangle_{\rm L}\hat{h}^{\mu\nu} := \nabla^2\hat{h}^{\mu\nu} + 2R^\mu{}_\alpha{}^\nu{}_\beta \hat{h}^{\alpha\beta} = 0\,, \tag{21}$$

where $\triangle_{\rm L}$ is the so-called Lichnerowicz operator.

As in the analysis of the original GL instability, we assume

$$\hat{h}^{\mu z} = 0\,, \quad \hat{h}^{zz} = 0\,. \tag{22}$$

Performing Fourier expansion along the symmetric spatial directions, we have

$$\hat{h}^{ab} = e^{i(\mu z + m\phi)}e^{\Omega v}h^{ab}(r,\theta), \tag{23}$$

where the subscripts a and b indicate the coordinates (v, r, θ, χ) and we have assumed that the frequency is pure imaginary as in Ref. [3]. Hereafter we concentrate on the $m = 0$ case. The assumption of a pure imaginary frequency is correct for the non-rotating case [1]. In our case, the validity of this assumption may be supported by the existence of such a solution. The equations of motion can be rewritten as

$$(\triangle_{\rm L}\hat{h})^{ab} = -\mu^2\hat{h}^{ab} + (\mathcal{L}\hat{h})^{ab} = 0 \;\Leftrightarrow\; -\mu^2\tilde{h}^{ab} + (\mathcal{L}\tilde{h})^{ab} = 0, \tag{24}$$

where

$$\tilde{h}^{ab} := e^{\Omega v}h^{ab} \tag{25}$$

and $\mathcal{L}$ is the Lichnerowicz operator in the Kerr spacetime. Gauge conditions are given by

$$\nabla_a \tilde{h}^{ab} = 0\,, \quad h^a{}_a = 0\,. \tag{26}$$

We solve these equations as an eigenvalue problem for μ with a fixed Ω.

As mentioned in Sec. I, we consider the slowly rotating case by introducing a small parameter ϵ. We specify the deviation from the non-rotating case by

$$M = \hat{M} + \epsilon^2 \hat{M}, \tag{27}$$

$$a = 2\epsilon \hat{M}. \tag{28}$$

In the limit $\epsilon \to 0$, the metric of a z =const. hyper-surface becomes the Schwarzschild metric. Since

$$\triangle = (r - 2\hat{M})(r - 2\hat{M}\epsilon^2), \tag{29}$$

we have

$$r_+ = 2\hat{M}, \quad r_- = \epsilon^2 r_+. \tag{30}$$

In this way, the coordinate radius of the horizon is unchanged by adding the rotation. This fact makes our analysis much simpler because the coordinate value at the horizon boundary does not depend on the order of ϵ.

Fixing the value of Ω, we expand $\mathcal{L}$, μ^2 and h as

$$\mathcal{L} = \mathcal{L}_0 + \epsilon \mathcal{L}_1 + \epsilon^2 \mathcal{L}_2 + \cdots\,, \tag{31}$$

$$\mu^2 = \mu_0^2 + \epsilon \mu_1^2 + \epsilon^2 \mu_2^2 + \cdots\,, \tag{32}$$

$$\tilde{h} = \tilde{h}_0 + \epsilon \tilde{h}_1 + \epsilon^2 \tilde{h}_2 + \cdots\,. \tag{33}$$

Since the value of μ should not depend on the direction of the rotation, we have $\mu_1^2 = 0$. Thus our goal is to get the value of μ_2^2. The equations of motion can be schematically described at each order of ϵ as

$$\epsilon^0 \;:\quad -\mu_0^2 \tilde{h}_0^{ab} + (\mathcal{L}_0 \tilde{h}_0)^{ab} = 0\,, \tag{34}$$

$$\epsilon^1 \;:\quad -\mu_0^2 \tilde{h}_1^{ab} + (\mathcal{L}_0 \tilde{h}_1)^{ab} + (\mathcal{L}_1 \tilde{h}_0)^{ab} = 0\,, \tag{35}$$

$$\epsilon^2 \;:\quad -\mu_0^2 \tilde{h}_2^{ab} + (\mathcal{L}_0 \tilde{h}_2)^{ab} + (\mathcal{L}_1 \tilde{h}_1)^{ab} + (\mathcal{L}_2 \tilde{h}_0)^{ab} - \mu_2^2 \tilde{h}_0^{ab} = 0\,. \tag{36}$$

In the following subsections, we find that there exist a single master variable that satisfies a second-order ordinary differential equation at each order of ϵ. The master variables must be carefully chosen so that the equation can be easily solved by numerical integration. An inappropriate choice of master variables may cause 0/0 terms in numerical integration as reported in Ref. [2]. For this reason, we treat the $\Omega = 0$ case separately from $\Omega \neq 0$ cases.

A. $\Omega = 0$ case

1. *Master Equations at order ϵ^0 and ϵ^1*

As the same as the original work by Gregory and Laflamme, we consider only s-wave perturbations at order ϵ^0. The metric perturbation can be written as

$$h_0^{ab} = \begin{pmatrix} h_0^{vv}(r) & h_0^{vr}(r) & 0 & 0 \\ h_0^{vr}(r) & h_0^{rr}(r) & 0 & 0 \\ 0 & 0 & h_0^{\theta\theta}(r) & 0 \\ 0 & 0 & 0 & \frac{h_0^{\theta\theta}(r)}{\sin^2\theta} \end{pmatrix} . \tag{37}$$

Here we introduce H_- defined by

$$H_- := \left(1 - \frac{r_+}{r}\right) h_0^{vv} - 2h_0^{vr} , \tag{38}$$

which is found to be a good master variable at order ϵ^0. Using the gauge conditions (26) and the equations of motion (34), we can derive the second-order ordinary differential equation for H_- ,

$$H_-'' = \left(\frac{3\mu_0^2 r^2 + 2}{\mu_0^2 r^3 + 2r - 2r_+} + \frac{1}{r_+ - r} - \frac{4}{r}\right) H_-' + \frac{\mu_0^2 r \left(\mu_0^2 r^3 + 4r - 8r_+\right)}{(r - r_+)\left(\mu_0^2 r^3 + 2r - 2r_+\right)} H_- . \tag{39}$$

All the components of the metric perturbation can be expressed in terms of H_-. The explicit expressions are given in Appendix B 1.

The metric perturbation at order ϵ^1 is induced by the perturbation at order ϵ^0 as given by Eq. (35). The only nontrivial induced components are found to be $h_1^{v\chi}$ and $h_1^{r\chi}$:

$$h_1^{ab} = \begin{pmatrix} 0 & 0 & 0 & h_1^{v\chi}(r) \\ 0 & 0 & 0 & h_1^{r\chi}(r) \\ 0 & 0 & 0 & 0 \\ h_1^{v\chi}(r) & h_1^{r\chi}(r) & 0 & 0 \end{pmatrix} . \tag{40}$$

Defining $f(r) := h_1^{v\chi}(r)$ as a master variable, we can derive the following equation:

$$f'' = \left(\frac{2}{r_+ - r} - \frac{2}{r}\right) f' + \frac{\left(\mu_0^2 r^3 - 2r_+\right)}{r^2(r - r_+)} f + S_1 ; \tag{41}$$

$$\begin{aligned} S_1 = & \frac{r_+\left(3rr_+^3\left(\mu_0^2 r^2 + 12\right) - 4r^2 r_+^2\left(\mu_0^2 r^2 + 7\right) + r^3 r_+\left(\mu_0^2 r^2 - 12\right) + 12r^4 - 12r_+^4\right)}{2r^3(r - r_+)^3\left(\mu_0^2 r^3 + 2r - 2r_+\right)} H_-' \\ & + \frac{r_+\left(r_+^3\left(10 - 3\mu_0^2 r^2\right) + 2rr_+^2\left(\mu_0^2 r^2 - 11\right) + r^2 r_+\left(7\mu_0^2 r^2 + 6\right) - 2\left(2\mu_0^2 r^5 + r^3\right)\right)}{r^3(r - r_+)^3\left(\mu_0^2 r^3 + 2r - 2r_+\right)} H_- . \end{aligned}$$

2. *Master equation at order ϵ^2*

Equations at order ϵ^2 are much more complicated than those to order ϵ^1. However, as we will see soon, we do not need to fully solve the perturbation equations to get the value of μ_2. Instead it is enough to focus on $\ell = 0$ modes in the spherical harmonics expansion. A minimal review of tensor and vector spherical harmonics is given in Appendix A. In Eq. (36), the first and second terms correspond to the homogeneous part of the coupled differential equations. The operator of this part is the same as that at order ϵ^0. This implies that different spherical harmonic modes of h_2 do not couple with each other within this homogeneous part. In addition, the final term proportional to μ_2^2 has only the $\ell = 0$ mode. Therefore, the value of μ_2^2 can be obtained by solving only the $\ell = 0$ mode of Eq. (36).

Extracting the $\ell = 0$ mode from Eq. (36), we obtain

$$-\mu_0^2 h_{2,0}^{ab} + (\mathcal{L}_0 h_{2,0})^{ab} + \Big[(\mathcal{L}_1 h_1)^{ab} + (\mathcal{L}_2 h_0)^{ab}\Big]_{\ell=0} - \mu_2^2 h_0^{ab} = 0\,, \tag{42}$$

where

$$h_{2,0}^{ab}(r) := \Big[h_2^{ab}(r)\Big]_{\ell=0}\,, \tag{43}$$

which can be expressed as

$$h_{2,0}^{ab} = \begin{pmatrix} h_2^{vv}(r) & h_2^{vr}(r) & 0 & 0 \\ h_2^{vr}(r) & h_2^{rr}(r) & 0 & 0 \\ 0 & 0 & h_2^{\theta\theta}(r) & 0 \\ 0 & 0 & 0 & \frac{h_2^{\theta\theta}(r)}{\sin^2\theta} \end{pmatrix}. \tag{44}$$

We also extract the $\ell = 0$ mode from the gauge conditions,

$$[\nabla_a h^a{}_b]_{l=0} = 0\,, \quad (h_{2,0})^a{}_a = 0\,. \tag{45}$$

Here in analogy with the case of order ϵ^0, we introduce the following master variable:

$$H_{2-} := \left(1 - \frac{r_+}{r}\right) h_2^{vv} - 2h_2^{vr}\,. \tag{46}$$

Then we obtain the master equation,

$$\begin{aligned}
H_{2-}'' = &\left(\frac{3\mu_0^2 r^2+2}{\mu_0^2 r^3+2r-2r_+}+\frac{1}{r_+-r}-\frac{4}{r}\right)H_{2-}' \\
&+\frac{\mu_0^2 r\left(\mu_0^2 r^3+4r-8r_+\right)}{(r-r_+)\left(\mu_0^2 r^3+2r-2r_+\right)}H_{2-}+S_2\,; \qquad (47)\\
S_2 = &\frac{4\mu_0^2 r r_+^2}{\mu_0^2 r^3+2r-2r_+}f'+\frac{4\mu_0^2 r_+^2(2r-r_+)}{(r-r_+)\left(\mu_0^2 r^3+2r-2r_+\right)}f \\
&+\frac{1}{3r^4(r-r_+)^2(\mu_0^2 r^3+2r-2r_+)^2}\Bigg(r_+\Big[-3\mu_0^4 r^8(r-r_+)(3r+r_+) \\
&+\mu_0^2 r^3(18r^5+4r^4 r_+-73r^3 r_+^2+14r^2 r_+^3+31 r r_+^4-12 r_+^5) \\
&+4r_+(r-r_+)(19r^4-33r^3 r_+-3r^2 r_+^2+17 r r_+^3-6r_+^4)\Big]+6\mu_2^2 r^6(2r-3r_+)(r-r_+)^2\Bigg)H_-' \\
&+\frac{1}{3r^4(r-r_+)^2(\mu_0^2 r^3+2r-2r_+)^2}\Bigg(3\mu_2^2 r^5(r-r_+)\Big[-4 r r_+(\mu_0^2 r^2+6) \\
&+r^2(\mu_0^4 r^4+4\mu_0^2 r^2+8)+16 r_+^2\Big]+r_+\Big[2r_+^5(3\mu_0^2 r^2-44)+r r_+^4(-3\mu_0^4 r^4+44\mu_0^2 r^2+208) \\
&-2r^2 r_+^3(2\mu_0^4 r^4+35\mu_0^2 r^2-24)+r^4 r_+(-2\mu_0^6 r^6+23\mu_0^4 r^4+84\mu_0^2 r^2+200) \\
&+3r^5(\mu_0^6 r^6-8\mu_0^4 r^4-16\mu_0^2 r^2-16)-r^3 r_+^2(\mu_0^6 r^6-26\mu_0^4 r^4+88\mu_0^2 r^2+320)\Big]\Bigg)H_-\,.
\end{aligned}$$

Explicit expressions for the metric components in terms of H_{2-} are given in Appendix B 3.

3. Asymptotic behavior

In order to solve the master equations, we need to specify the boundary conditions. In the limit $r \to \infty$, Eq. (39) can be approximated as

$$H_-'' + \frac{2}{r}H_-' - \mu_0^2 H_- = 0\,. \qquad (48)$$

Hence the general solution at $r \to \infty$ is given by

$$H_- = C_1\frac{e^{-\mu_0 r}}{r} + C_2\frac{e^{\mu_0 r}}{r}\,, \qquad (49)$$

where we may assume $\mu_0 > 0$ without loss of generality. Then the regularity requires $C_2 = 0$. Thus H_- decays exponentially at infinity, and so do all the components of the metric at order ϵ^0. At order ϵ, the homogeneous solution of the master equation is found to have the similar asymptotic behavior. Since the source term is given in terms of H_- and H_-', this implies that the master variable $f(r)$, and hence all the components of the metric perturbation at order ϵ also decay exponentially at infinity. The same is true for the master equation at order ϵ^2 since it has the same form as the master equation at order ϵ^0. Thus all the components of the metric perturbations to order ϵ^2 must decay exponentially at the infinity.

4. Near the horizon

From the regularity at the horizon, we find $H_- = \mathcal{O}((r-r_+)^0)$, $f = \mathcal{O}((r-r_+)^0)$ and $H_{2-} = \mathcal{O}((r-r_+)^0)$. Expanding the zeroth and first order master equations near the horizon, we obtain the following asymptotic solutions for H_- and f:

$$\begin{aligned} H_- &= \alpha\left(1+\left(\mu_0^2 r_+ - \frac{4}{r_+}\right)(r-r_+) + \left(\frac{\mu_0^4 r_+^2}{4} - 2\mu_0^2 + \frac{8}{r_+^2}\right)(r-r_+)^2\right) \\ &\quad +\mathcal{O}((r-r_+)^3), \end{aligned} \tag{50}$$

$$f = \beta - \frac{(r-r_+)\left(-6\beta\mu_0^2 r_+^3 + 12\beta r_+ + \mu_0^4 r_+^4 - 19\mu_0^2 r_+^2 + 94\right)}{12 r_+^2} + \mathcal{O}\left((r-r_+)^2\right), \tag{51}$$

where α and β are constants to be determined. Without loss of generality, we may set $\alpha = 1$ because Eq. (39) is a homogeneous linear equation. The eigenvalue μ_0^2 is determined so that H_- decays exponentially at infinity, and the constant β is determined so that f decays exponentially at infinity.

Since the zeroth order equation (39) and the second order equation (47) have the same form except for the presence of the source term S_2 for the latter, one can add the zeroth order solution H_- to any particular solution H_{2-} of Eq. (47). Using this freedom, we can set $H_{2-}(r_+) = 0$ by adding a term proportional to H_- appropriately. Then expanding Eq. (47) near the horizon, we find

$$H_{2-} = (r-r_+)\left(4\beta + r_+\left(\frac{\mu_0^2}{3} + \mu_2^2\right) - \frac{7}{r_+}\right) + \mathcal{O}\left((r-r_+)^2\right). \tag{52}$$

As before, the value of μ_2^2 is determined so that H_{2-} exponentially decays at infinity.

B. $\Omega \neq 0$ case

In the $\Omega \neq 0$ case if we would use the same master variables as those in the $\Omega = 0$ case, we would suffer from 0/0 terms in the master equations at order ϵ^0 and ϵ^2, as pointed out in Ref. [2]. To avoid numerical errors due to these 0/0 terms, as in Ref. [2], we rewrite each master equation at order ϵ^0 and ϵ^2 as two coupled first-order differential equations by introducing an additional variable.

At order ϵ^0, we define a function H as

$$H := -h_0^{vr} + \frac{1}{1-\frac{r_+}{r}} h_0^{rr}. \tag{53}$$

Then, we obtain the following equations for H_- and H:

$$\begin{aligned} H'_- &= \frac{\left(\mu_0^4 r^3 + 2\mu_0^2(r-r_+) - 4\Omega^2 r\right)}{\Omega\left(\mu_0^2 r^3 + r_+\right)} H \\ &+ \frac{\left(\mu_0^2 r^3(3r_+ - 2r(\Omega r+1)) + 4\Omega^2 r^4 - 2rr_+(\Omega r+3) + 6r_+^2\right)}{2r(r-r_+)\left(\mu_0^2 r^3 + r_+\right)} H_- \,, \end{aligned} \tag{54}$$

$$\begin{aligned} H' &= \frac{\Omega r\left(4\mu_0^2 r^3(r-r_+) + 4\Omega^2 r^4 - r_+^2\right)}{4(r-r_+)^2\left(\mu_0^2 r^3 + r_+\right)} H_- \\ &- \frac{\left(\mu_0^2 r^3(2r(\Omega r+1) + r_+) + 4\Omega^2 r^4 + 2rr_+(\Omega r+2) - 2r_+^2\right)}{2r(r-r_+)\left(\mu_0^2 r^3 + r_+\right)} H \,. \end{aligned} \tag{55}$$

These equations are free from 0/0 terms. The components of the metric perturbation are expressed in terms of H_- and H as shown in Appendix C.

At order ϵ^1, we can use the same master variable as the one in the $\Omega = 0$ case, $f := h_1^{v\chi}$. It satisfies

$$\begin{aligned} f'' &= \left(\frac{\mu_0^2}{\mu_0^2 r + 4\Omega} + \frac{2(\Omega r+1)}{r_+ - r} - \frac{3}{r}\right) f' + \frac{\mu_0^4 r^4 - 2\mu_0^2 r r_+ - 24\Omega^2 r^2 - 12\Omega r_+}{r^2(r-r_+)\left(\mu_0^2 r + 4\Omega\right)} f \\ &\quad + A_H(r; r_+, \mu_0, \Omega) H + A_{H_-}(r; r_+, \mu_0, \Omega) H_- \,, \end{aligned} \tag{56}$$

where explicit expressions of $A_H(r; r_+, \mu_0, \Omega)$ and $A_{H_-}(r; r_+, \mu_0, \Omega)$ are given in Appendix C. The other component $h_1^{r\chi}$ of the metric perturbation is also expressed in terms of f in Appendix C.

At order ϵ^2, we define H_2 as

$$H_2 := -h_2^{vr} + \frac{1}{1 - \frac{r_+}{r}} h_2^{rr}. \tag{57}$$

Then, we can derive

$$\begin{aligned} H'_{2-} &= \frac{(4\Omega^2 r^4 + \mu_0^2(3r_+ - 2r(\Omega r+1))r^3 - 2(\Omega r+3)r_+ r + 6r_+^2)}{2r(r-r_+)(\mu_0^2 r^3 + r_+)} H_{2-} \\ &\quad + \frac{(r^3\mu_0^4 + 2(r-r_+)\mu_0^2 - 4\Omega^2 r)}{\Omega(\mu_0^2 r^3 + r_+)} H_2 \\ &\quad + \frac{4(r\mu_0^2 + 2\Omega)(r-r_+)r_+^2}{r(r\mu_0^2 + 4\Omega)(\mu_0^2 r^3 + r_+)} f' + \frac{(8r(r\mu_0^2 + \Omega(\mu_0^2 r^2 + 2\Omega r + 4))r_+^2 - 4(r\mu_0^2 + 6\Omega)r_+^3)}{r^2(r\mu_0^2 + 4\Omega)(\mu_0^2 r^3 + r_+)} f \\ &\quad + B_H(r; r_+, \mu_0, \mu_2, \Omega) H + B_{H_-}(r; r_+, \mu_0, \mu_2, \Omega) H_-, \end{aligned} \tag{58}$$

$$\begin{aligned} H'_2 &= \frac{\Omega r(4\Omega^2 r^4 + 4\mu_0^2(r-r_+)r^3 - r_+^2)}{4(r-r_+)^2(\mu_0^2 r^3 + r_+)} H_{2-} \\ &\quad - \frac{(4\Omega^2 r^4 + \mu_0^2(2r(\Omega r+1) + r_+)r^3 + 2(\Omega r+2)r_+ r - 2r_+^2)}{2r(r-r_+)(\mu_0^2 r^3 + r_+)} H_2 \\ &\quad + \frac{2\Omega(2(r\mu_0^2 + \Omega)r^2 + r_+)r_+^2}{r(r\mu_0^2 + 4\Omega)(\mu_0^2 r^3 + r_+)} f' + \frac{2\Omega(2(r\mu_0^2 + 2\Omega(r(r\mu_0^2 + \Omega) + 2))r^3 - 4\Omega r_+ r^2 + r_+^2)r_+^2}{r^2(r\mu_0^2 + 4\Omega)(r-r_+)(\mu_0^2 r^3 + r_+)} f \\ &\quad + C_H(r; r_+, \mu_0, \mu_2, \Omega) H + C_{H_-}(r; r_+, \mu_0, \mu_2, \Omega) H_- \,, \end{aligned} \tag{59}$$

where explicit expressions of $B_H(r; r_+, \mu_0, \mu_2, \Omega)$, $B_{H_-}(r; r_+, \mu_0, \mu_2, \Omega)$, $C_H(r; r_+, \mu_0, \mu_2, \Omega)$ and $C_H(r; r_+, \mu_0, \mu_2, \Omega)$ are given in Appendix C. The components of the metric perturbation are also shown in Appendix C.

At infinity, all these variables must exponentially decay. We can derive asymptotic solutions near the horizon as in the $\Omega = 0$ case. We describe these asymptotic solutions near the horizon in Appendix C.

IV. RESULTS AND DISCUSSION

Since a second-order ordinary differential equation can be regarded as a set of two coupled first-order differential equations, we have six coupled first-order ordinary differential equations for each value of Ω. We have to determine the values of μ_0, β and μ_2 so that the metric perturbation is regular at horizon and decays exponentially at infinity. It can be done numerically by a shooting method. We have solved the ordinary differential equations from the horizon to infinity using the 4-th order Runge-Kutta method and searched for appropriate values of μ_0, β and μ_2 for each value of Ω.

Here, we show the numerical results by fixing the mass parameter M of the Kerr black hole at the 4-dimensional section, although the numerical integration is done by fixing the value of r_+. That is, we plot ΩM as a function of μM, where $\mu = \sqrt{\mu_0^2 + \mu_2^2 \epsilon^2}$, in Fig. 1, where

$$\Omega M \;=\; \Omega \hat{M}(1+\epsilon^2)\,, \tag{60}$$

$$\mu M \;=\; \sqrt{\mu_0^2 + \mu_2^2\epsilon^2}\,\hat{M}(1+\epsilon^2) = \mu_0 \hat{M}\left(1 + \left(1 + \frac{1}{2}\frac{\mu_2^2}{\mu_0^2}\right)\epsilon^2\right) + \mathcal{O}(\epsilon^3)\,. \tag{61}$$

It is clear from the figure that both the critical wavelength and the growth time-scale of the instability become shorter as the rotation is increased. In other words, rotation makes the black string system more unstable. This result is consistent with previous works in similar situations as listed in Sec. I.

Although our results are consistent with all the previous results, an intuitive understanding of our results seems rather unclear. If we fix the mass parameter of the Kerr black hole in the 4-dimensional section, the horizon area decreases as the angular momentum is increased. In this sense, the typical time-scale or length-scale becomes shorter. This is consistent with our and previous results, and this was in fact what we obtained from thermodynamic arguments. However, in contrast, if we focus on the surface gravity, it decreases as the angular momentum is increased. In this respect, the typical time-scale or length-scale may become longer. Actually, when we add a charge to a neutral black brane solution, it is known that the GL instability gets weaker [2]. This is totally opposite to the behavior we obtained. Apparently we need a better physical picture of the GL instability with rotation.

Acknowledgements

We thank T. Tanaka, G. Kang for helpful discussions and comments. We made much progress during the APCTP Joint Focus Program: Frontiers of Black Hole Physics (December 2010). We would like to thank all the participants of this workshop for fruitful discussions. C.Y. is supported by JSPS Grant-in-Aid for Creative Scientific Research No. 19GS0219 and JSPS Grant-in-Aid for



FIG. 1: The growth rate of the GL instability Ω as a function of the eigenvalue $\mu(=\sqrt{\mu_0^2+\epsilon^2\mu_2^2})$. Plotted are the case of $\epsilon=0$(no rotation) and $\epsilon=0.2$.

Scientific Research (C) No. 21540276. This work was also supported in part by Korea Institute for Advanced Study under the KIAS Scholar program, by the Grant-in-Aid for the Global COE Program "The Next Generation of Physics, Spun from Universality and Emergence" from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan, by JSPS Grant-in-Aid for Scientific Research (A) No. 21244033, and by JSPS Grant-in-Aid for Creative Scientific Research No. 19GS0219.

Appendix A: Spherical Harmonics

Full expressions for the tensor spherical harmonics can be found in Refs. [16–18]. Here, we give a minimum review necessary for the present work. We represent tensors by using the components of the coordinate system (v,r,θ,ϕ), where v is an advanced null coordinate. First, we introduce the vector spherical harmonics. We decompose any covariant vector field $\boldsymbol{v}$ as

$$\boldsymbol{v}=\sum_{l,m}\left[A_{lm}\boldsymbol{a}_{lm}+B_{lm}\boldsymbol{b}_{lm}+C_{lm}\boldsymbol{c}_{lm}+D_{lm}\boldsymbol{d}_{lm}\right], \tag{A1}$$

where A, B, C, D are the coefficients for bases $\boldsymbol{a},\ \boldsymbol{b},\ \boldsymbol{c},\ \boldsymbol{d}$ given by

$$\begin{aligned}
\boldsymbol{a}_{lm} &= -i(Y_{lm}, -Y_{lm}, 0, 0)\,, && \text{(A2)}\\
\boldsymbol{b}_{lm} &= (0, Y_{lm}, 0, 0)\,, && \text{(A3)}\\
\boldsymbol{c}_{lm} &= \frac{1}{\sqrt{l(l+1)}}(0, 0, \partial_\theta Y_{lm}, \partial_\phi Y_{lm})\,, && \text{(A4)}\\
\boldsymbol{d}_{lm} &= \frac{1}{\sqrt{l(l+1)}}(0, 0, \frac{1}{\sin\theta}\partial_\phi Y_{lm}, -\sin\theta\partial_\theta Y_{lm})\,. && \text{(A5)}
\end{aligned}$$

These bases are orthonormal in the inner product

$$(\boldsymbol{v}, \boldsymbol{u}) := \int \eta^{ab} v_a u_b \mathrm{d}\Omega, \tag{A6}$$

where η is the Minkowski metric.

We focus on the $l = m = 0$ mode. In this case, we have the following two non-zero bases:

$$\boldsymbol{a} := \boldsymbol{a}_{00} = \frac{-i}{2\sqrt{\pi}}(1,\ -1,\ 0,\ 0)\ ,\quad \boldsymbol{b} := \boldsymbol{b}_{00} = \frac{1}{2\sqrt{\pi}}(0,\ 1,\ 0,\ 0). \tag{A7}$$

Thus, we can extract the $l = m = 0$ mode from a covariant vector $\boldsymbol{v}$ as

$$[\boldsymbol{v}]_{l=0} = (\boldsymbol{a}, \boldsymbol{v})\boldsymbol{a} + (\boldsymbol{b}, \boldsymbol{v})\boldsymbol{b}. \tag{A8}$$

In the same way as in the vector spherical harmonics, we can decompose a rank-2 covariant tensor $\mathbf{v}$ as follows:

$$\begin{aligned}
v &= \sum_{l,m}\Big[A_{lm}\mathbf{a}_{lm} + B_{lm}\mathbf{b}_{lm} + C_{lm}\mathbf{c}_{lm} + D_{lm}\mathbf{d}_{lm} + E_{lm}\mathbf{e}_{lm} + F_{lm}\mathbf{f}_{lm}\\
&\quad + G_{lm}\mathbf{g}_{lm} + H_{lm}\mathbf{h}_{lm} + I_{lm}\mathbf{i}_{lm} + J_{lm}\mathbf{j}_{lm}\Big], && \text{(A9)}
\end{aligned}$$

where $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$, $\mathbf{d}$, $\mathbf{e}$, $\mathbf{f}$, $\mathbf{g}$, $\mathbf{h}$, $\mathbf{i}$, $\mathbf{j}$ are bases of the tensor spherical harmonics and $A,\ B,\ C, \cdots,\ I,\ J$ are the coefficients for these bases. Explicit forms of the bases are given in Ref. [18]. These bases are orthogonal in the inner product

$$< \mathbf{v},\ \mathbf{u} >:= \int \eta^{ac}\eta^{bd} v_{ab} u_{cd} \mathrm{d}\Omega. \tag{A10}$$

For $m = l = 0$, we have the following four non-zero bases:

$$\mathbf{a} := \mathbf{a}_{0,0} = \frac{1}{2\sqrt{\pi}} \begin{pmatrix} 1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \tag{A11}$$

$$\mathbf{b} := \mathbf{b}_{0,0} = \frac{i}{2\sqrt{2\pi}} \begin{pmatrix} 0 & -1 & 0 & 0 \\ -1 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \tag{A12}$$

$$\mathbf{c} := \mathbf{c}_{0,0} = \frac{1}{2\sqrt{\pi}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \tag{A13}$$

$$\mathbf{d} := \mathbf{d}_{0,0} = \frac{r^2}{2\sqrt{2\pi}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & \sin^2\theta \end{pmatrix}. \tag{A14}$$

We can extract the $l = m = 0$ mode from $\mathbf{v}$ as

$$[\mathbf{v}]_{l=0} = < \mathbf{a}, \mathbf{v} > \mathbf{a} + < \mathbf{b}, \mathbf{v} > \mathbf{b} + < \mathbf{c}, \mathbf{v} > \mathbf{c} + < \mathbf{d}, \mathbf{v} > \mathbf{d}. \tag{A15}$$

Appendix B: metric perturbation for $\Omega = 0$

Here we give explicit expressions for the components of the metric perturbation at each order of ϵ in the case $\Omega = 0$. For notational simplicity, ϵ is set to unity in the following expressions.

1. Metric perturbation at order ϵ^0

$$h_0^{vv} = \frac{r^2(2r-3r_+)}{(r-r_+)\left(\mu_0^2 r^3 + 2r - 2r_+\right)} H_-' + \frac{r\left(\mu_0^2 r^3 + 4r - 8r_+\right)}{(r-r_+)\left(\mu_0^2 r^3 + 2r - 2r_+\right)} H_-\,, \tag{B1}$$

$$h_0^{vr} = \frac{r(2r-3r_+)}{2\mu_0^2 r^3 + 4r - 4r_+} H_-' + \frac{r - 3r_+}{\mu_0^2 r^3 + 2r - 2r_+} H_-\,, \tag{B2}$$

$$h_0^{rr} = \frac{(2r-3r_+)(r-r_+)}{2\mu_0^2 r^3 + 4r - 4r_+} H_-' + \frac{(r-3r_+)(r-r_+)}{r\left(\mu_0^2 r^3 + 2r - 2r_+\right)} H_-\,, \tag{B3}$$

$$h_0^{\theta\theta} = \frac{1}{2r^2} H_-\,. \tag{B4}$$

2. Metric perturbation at order ϵ^1

$$h_1^{r\chi} = \frac{r_+(2r-3r_+)}{2\mu_0^2 r^4+4r^2-4rr_+}H'_- + \frac{r_+(r-3r_+)}{r^2\left(\mu_0^2 r^3+2r-2r_+\right)}H_-. \tag{B5}$$

3. Metric perturbation at order ϵ^2

$$\begin{aligned} h_2^{rr} \;=\; & \frac{(2r-3r_+)(r-r_+)}{2\mu_0^2 r^3+4r-4r_+}H'_{2-} + \frac{(r-3r_+)(r-r_+)}{r\left(\mu_0^2 r^3+2r-2r_+\right)}H_{2-} \\ & + \frac{2r_+^2(r-r_+)^2}{r\left(\mu_0^2 r^3+2r-2r_+\right)}h_1^{v\chi\prime} + \frac{2r_+^2(r-r_+)(2r-r_+)}{r^2\left(\mu_0^2 r^3+2r-2r_+\right)}h_1^{v\chi} \\ & + \Bigg(r_+\Big(-21\mu_0^2 r^7+31\mu_0^2 r^6 r_+ + 2r^5(\mu_0^2 r_+^2-3) + \\ & r^4 r_+(40-9\mu_0^2 r_+^2) - r^3 r_+^2(3\mu_0^2 r_+^2+59) + 4r^2 r_+^3 + 9rr_+^4 + 6r_+^5\Big) \\ & -3\mu_2^2 r^6(2r-3r_+)(r-r_+)\Bigg)\frac{H'_-}{6r^3(\mu_0^2 r^3+2r-2r_+)^2} \\ & + \Bigg(r_+\Big(-36\mu_0^2 r^7+79\mu_0^2 r^6 r_+ - 4r^5(4\mu_0^2 r_+^2+9) \\ & +8r^4 r_+(14-3\mu_0^2 r_+^2) + r^3 r_+^2(3\mu_0^2 r_+^2-176) + 34r^2 r_+^3 + 48rr_+^4 - 6r_+^5\Big) \\ & -6\mu_2^2 r^6(r-3r_+)(r-r_+)\Bigg)\frac{H_-}{6r^4(\mu_0^2 r^3+2r-2r_+)^2}, \end{aligned} \tag{B6}$$

$$\begin{aligned} h_2^{vv} \;=\; & \frac{r^2(2r-3r_+)}{(r-r_+)\left(\mu_0^2 r^3+2r-2r_+\right)}H'_{2-} + \frac{r\left(\mu_0^2 r^3+4r-8r_+\right)}{(r-r_+)\left(\mu_0^2 r^3+2r-2r_+\right)}H_{2-} \\ & + \frac{4rr_+^2}{\mu_0^2 r^3+2r-2r_+}h_1^{v\chi\prime} + \frac{4r_+^2(2r-r_+)}{(r-r_+)\left(\mu_0^2 r^3+2r-2r_+\right)}h_1^{v\chi} \\ & + \Bigg(r_+\Big(-15\mu_0^2 r^7 - 3rr_+^4(\mu_0^2 r^2+3) \\ & +2r^4 r_+(11\mu_0^2 r^2+5) - r^3 r_+^2(4\mu_0^2 r^2+53) + 6r^5 + 34r^2 r_+^3 + 6r_+^5\Big) \\ & -3\mu_2^2 r^6(2r-3r_+)(r-r_+)\Bigg)\frac{H'_-}{3r(r-r_+)^2(\mu_0^2 r^3+2r-2r_+)^2} \\ & + \Bigg(r_+\Big(3rr_+^4(\mu_0^2 r^2+4) + 2r^2 r_+^3(41-3\mu_0^2 r^2) \\ & -6r^5(5\mu_0^2 r^2+4) + r^4 r_+(61\mu_0^2 r^2+64) - 2r^3 r_+^2(11\mu_0^2 r^2+76) - 6r_+^5\Big) \\ & -6\mu_2^2 r^6(r-3r_+)(r-r_+)\Bigg)\frac{H_-}{3r^2(r-r_+)^2(\mu_0^2 r^3+2r-2r_+)^2}, \end{aligned} \tag{B7}$$

$$\begin{aligned} h_2^{vr} \;=\; & \frac{r(2r-3r_+)}{2\mu_0^2 r^3+4r-4r_+}H'_{2-} + \frac{(r-3r_+)}{\mu_0^2 r^3+2r-2r_+}H_{2-} \\ & + \frac{2r_+^2(r-r_+)}{\mu_0^2 r^3+2r-2r_+}h_1^{v\chi\prime} + \frac{2r_+^2(2r-r_+)}{\mu_0^2 r^4+2r^2-2rr_+}h_1^{v\chi} \\ & + \Bigg(r_+\Big(-15\mu_0^2 r^7 - 3rr_+^4(\mu_0^2r^2+3) + 2r^4 r_+(11\mu_0^2 r^2+5) - r^3 r_+^2(4\mu_0^2 r^2+53) + 6r^5 \\ & +34r^2 r_+^3 + 6r_+^5\Big) - 3\mu_2^2 r^6(2r-3r_+)(r-r_+)\Bigg)\frac{H'_-}{6r^2(r-r_+)(\mu_0^2 r^3+2r-2r_+)^2} \\ & + \Bigg(r_+\Big(3rr_+^4(\mu_0^2 r^2+4) + 2r^2 r_+^3(41-3\mu_0^2 r^2) \\ & -6r^5(5\mu_0^2 r^2+4) + r^4 r_+(61\mu_0^2 r^2+64) - 2r^3 r_+^2(11\mu_0^2 r^2+76) - 6r_+^5\Big) \\ & -6\mu_2^2 r^6(r-3r_+)(r-r_+)\Bigg)\frac{H_-}{6r^3(r-r_+)(\mu_0^2 r^3+2r-2r_+)^2}, \end{aligned} \tag{B8}$$

$$\begin{aligned} h_2^{\theta\theta} \;=\; & \frac{1}{2r^2}H_{2-} + \frac{2r_+^2}{3r^3}h_1^{v\chi} - \frac{r_+(2r-3r_+)(3r^2-2rr_++r_+^2)}{6r^3(r-r_+)(\mu_0^2 r^3+2r-2r_+)}H'_- \\ & + r_+\Big(-3\mu_0^2 r^6 - 2\mu_0^2 r^5 r_+ + 2r^4\left(\mu_0^2 r_+^2-6\right) + r^3 r_+(\mu_0^2 r_+^2+24) \\ & -6r^2 r_+^2 + 4rr_+^3 - 2r_+^4\Big)\frac{H_-}{6r^5(r-r_+)\left(\mu_0^2 r^3+2r-2r_+\right)}\,. \end{aligned} \tag{B9}$$

Appendix C: expressions for $\Omega \neq 0$

1. Metric perturbation at order ϵ^0

$$\begin{aligned} h_0^{vv} \;=\; & \frac{r\left(2\mu_0^2 r^3(r-r_+) + 4\Omega^2 r^4 + r_+(r_+-2r)\right)}{2(r-r_+)^2\left(\mu_0^2 r^3+r_+\right)}H_- \\ & - \frac{r\left(\mu_0^2 r(2r(\Omega r-1)+3r_+) + 2\Omega\left(2\Omega r^2+r_+\right)\right)}{\Omega(r-r_+)\left(\mu_0^2 r^3+r_+\right)}H, \end{aligned} \tag{C1}$$

$$h_0^{vr} \;=\; \frac{4\Omega^2 r^4 + r_+(3r_+-4r)}{4(r-r_+)(\mu_0^2 r^3+r_+)}H_- - \frac{\mu_0^2 r(2r(\Omega r-1)+3r_+) + 2\Omega(2\Omega r^2+r_+)}{2\Omega(\mu_0^2 r^3+r_+)}H, \tag{C2}$$

$$h_0^{rr} \;=\; \frac{(r-r_+)\left(\mu_0^2(2r-3r_+) - 4\Omega^2 r\right)}{2\Omega\left(\mu_0^2 r^3+r_+\right)}H + \frac{\left(4\Omega^2 r^4 - 4rr_+ + 3r_+^2\right)}{4\mu_0^2 r^4+4rr_+}H_-, \tag{C3}$$

$$h_0^{\theta\theta} \;=\; \frac{1}{2r^2}H_-\,. \tag{C4}$$

2. Metric perturbation at order ϵ^1

$$
\begin{aligned}
h_1^{r\chi} \;=\;& \frac{r_+}{2\Omega r^2(r-r_+)(\mu_0^2 r+4\Omega)(\mu_0^2 r^3+r_+)}\Big(\mu_0^4 r^2(2r-3r_+)(r-r_+) \\
&+2\mu_0^2\Omega\Big(2\Omega r^4+r_+\left(-4r^2+8rr_+-3r_+^2\right)\Big)+4\Omega^2 r_+(2\Omega r+1)(2r-r_+)\Big)H \\
&+\frac{r_+}{4r^3(r-r_+)^2\left(\mu_0^2 r+4\Omega\right)\left(\mu_0^2 r^3+r_+\right)}\Big(\mu_0^2 r^2(r-r_+)(4\Omega r^3-6\Omega r r_+^2-4rr_++3r_+^2) \\
&+2\Omega r_+\Big(-2r^3\big(\Omega r(4\Omega r+1)-1\big)+2r^2 r_+\big(\Omega r(2\Omega r+1)+3\big)-13rr_+^2+6r_+^3\Big)\Big)H_- \\
&+\frac{\Omega(r_+-r)}{\mu_0^2 r+4\Omega}h_1^{v\chi\prime}-\frac{\Omega\left(2\Omega r^2+r_+\right)}{r\left(\mu_0^2 r+4\Omega\right)}h_1^{v\chi}.
\end{aligned}
\tag{C5}
$$

3. Metric perturbation at order ϵ^2

$$
\begin{aligned}
h_2^{rr} \;=\;& \frac{2(r-r_+)\Big(2r\big(r\mu_0^2+\Omega(\mu_0^2r^2+2\Omega r+4)\big)-(r\mu_0^2+6\Omega)r_+\Big)r_+^2}{r^2(r\mu_0^2+4\Omega)(\mu_0^2r^3+r_+)}h_1^{v\chi} \\
&+\frac{2(r\mu_0^2+2\Omega)(r-r_+)^2r_+^2}{r(r\mu_0^2+4\Omega)(\mu_0^2r^3+r_+)}h_1^{v\chi\prime} \\
&+\frac{4\Omega^2r^4-4r_+r+3r_+^2}{4\mu_0^2r^4+4r_+r}H_{2-}-\frac{(r-r_+)\Big((3r_+-2r)\mu_0^2+4\Omega^2r\Big)}{2\Omega\left(\mu_0^2r^3+r_+\right)}H_2 \\
&-\frac{1}{24\Omega r^4(r\mu_0^2+4\Omega)(r-r_+)(\mu_0^2r^3+r_+)^2}\Bigg(6\mu_2^2\Omega(r\mu_0^2+4\Omega)(r-r_+)\big(4\Omega^2r^4+r_+(3r_+-4r)\big)r^6 \\
&+r_+\Bigg[r^3(r-r_+)\Big(24\Omega\big(\Omega r(\Omega r-1)+1\big)r^5+4\big(\Omega r(\Omega r(2\Omega r+3)-15)+6\big)r_+r^3 \\
&-2\big(\Omega r(6\Omega r-17)+15\big)r_+^2r^2-3(2\Omega r+9)r_+^3r+3(9-16\Omega r)r_+^4\Big)\mu_0^4 \\
&+\Omega r\Big(24\Omega(\Omega r(2\Omega r(\Omega r+3)-3)+4)r^7+16(\Omega r(\Omega r(2\Omega r(\Omega r-2)+3)-24)+6)r_+r^5 \\
&+2(2\Omega r(4\Omega r(\Omega r(\Omega r-5)-8)+79)-105)r_+^2r^4-4(\Omega r(19\Omega r+16)-42)r_+^3r^3 \\
&+3(4\Omega r(7\Omega r-1)+39)r_+^4r^2+3(16\Omega r-81)r_+^5r+102r_+^6\Big)\mu_0^2+4\Omega^2(24\Omega^2(2\Omega r(\Omega r+1)+1)r^8 \\
&+16\Omega(2\Omega^3r^3-6\Omega r-3)r_+r^6+2(2\Omega r(4\Omega r(\Omega r(\Omega r-3)-2)+9)+3)r_+^2r^4 \\
&+4(\Omega r(6-7\Omega r)+9)r_+^3r^3+3(2\Omega r-1)(6\Omega r+1)r_+^4r^2-51r_+^5r+30r_+^6)\Bigg]\Bigg)H_- \\
&+\frac{1}{12\Omega^2r^3(r\mu_0^2+4\Omega)(\mu_0^2r^3+r_+)^2}\Bigg(6\mu_2^2\Omega(r\mu_0^2+4\Omega)(r-r_+)\big(4\Omega^2r^4+(2r-3r_+)r_+\big)r^3 \\
&+r_+\Bigg[-3r^3(r-r_+)\Big(10\Omega r^4-2(\Omega r_++2)r^3+2r_+(4-3\Omega r_+)r^2+r_+^2(3-2\Omega r_+)r-9r_+^3\Big)\mu_0^6 \\
&+\Omega r\Big(12(2\Omega^3r^3-14\Omega r+3)r^5+4(\Omega r(\Omega r(6\Omega r+5)+48)-30)r_+r^4 \\
&+2(\Omega r(2\Omega r+21)+50)r_+^2r^3+(83-102\Omega r)r_+^3r^2-3(8\Omega r+59)r_+^4r+48r_+^5\Big)\mu_0^4 \\
&+2\Omega^2\Big(12(\Omega r(2\Omega r(3\Omega r+1)-7)-2)r^5+4(\Omega r(\Omega r(16\Omega r+13)+18)+12)r_+r^4 \\
&+2\big(\Omega r(2\Omega r(2\Omega r-1)+13)-14\big)r_+^2r^3-(176\Omega r+5)r_+^3r^2+3(10\Omega r-27)r_+^4r+24r_+^5\Big)\mu_0^2 \\
&+8\Omega^3\Big(12\Omega(2\Omega r(\Omega r+1)+1)r^5+4\Omega(\Omega r(4\Omega r+3)-6)r_+r^4 \\
&+2\big(\Omega r(2\Omega r(2\Omega r-3)-5)-6\big)r_+^2r^2-(20\Omega r+3)r_+^3r+3(2\Omega r-1)r_+^4\Big)\Bigg]\Bigg)H, \qquad \text{(C6)}
\end{aligned}
$$

$$
\begin{aligned}
h_2^{vv} \;=\;& \frac{4r(r\mu_0^2+2\Omega)r_+^2}{(r\mu_0^2+4\Omega)(\mu_0^2r^3+r_+)}h_1^{v\chi\prime}+\frac{8r(r\mu_0^2+\Omega(\mu_0^2r^2+2\Omega r+4))r_+^2-4(r\mu_0^2+6\Omega)r_+^3}{(r\mu_0^2+4\Omega)(r-r_+)(\mu_0^2r^3+r_+)}h_1^{v\chi}\\
&+\frac{r(4\Omega^2r^4+2\mu_0^2(r-r_+)r^3+r_+(r_+-2r))}{2(r-r_+)^2(\mu_0^2r^3+r_+)}H_{2-}\\
&+\frac{r}{r-r_+}\left(\frac{r(\mu_0^2(2r-3r_+)-4\Omega^2r)}{\Omega(\mu_0^2r^3+r_+)}-2\right)H_2\\
&-\frac{1}{12\Omega r^2(r\mu_0^2+4\Omega)(r-r_+)^3(\mu_0^2r^3+r_+)^2}\\
&\times\Bigg(6\mu_2^2\Omega(r\mu_0^2+4\Omega)(r-r_+)(4\Omega^2r^4+r_+(3r_+-4r))r^6\\
&+r_+\Bigg[r^3(r-r_+)\Big(24\Omega(\Omega r(\Omega r-1)+1)r^5+4(\Omega r(\Omega r(2\Omega r+3)-15)+6)r_+r^3\\
&-2(\Omega r(6\Omega r-17)+15)r_+^2r^2-3(2\Omega r+9)r_+^3r+3(9-16\Omega r)r_+^4\Big)\mu_0^4\\
&+\Omega r(24\Omega(\Omega r(2\Omega r(\Omega r+3)-3)+4)r^7+16(\Omega r(\Omega r(2\Omega r(\Omega r-2)+3)-24)+6)r_+r^5\\
&+2(2\Omega r(4\Omega r(\Omega r(\Omega r-5)-8)+79)-105)r_+^2r^4-4(\Omega r(19\Omega r+16)-42)r_+^3r^3\\
&+3(4\Omega r(7\Omega r-1)+39)r_+^4r^2+3(16\Omega r-81)r_+^5r+102r_+^6)\mu_0^2+4\Omega^2(24\Omega^2(2\Omega r(\Omega r+1)+1)r^8\\
&+16\Omega(2\Omega^3r^3-6\Omega r-3)r_+r^6+2(2\Omega r(4\Omega r(\Omega r(\Omega r-3)-2)+9)+3)r_+^2r^4\\
&+4(\Omega r(6-7\Omega r)+9)r_+^3r^3+3(2\Omega r-1)(6\Omega r+1)r_+^4r^2-51r_+^5r+30r_+^6)\Bigg]\Bigg)H_-\\
&+\frac{1}{6\Omega^2r(r\mu_0^2+4\Omega)(r-r_+)^2(\mu_0^2r^3+r_+)^2}\\
&\times\Bigg(6\mu_2^2\Omega(r\mu_0^2+4\Omega)(r-r_+)(4\Omega^2r^4+(2r-3r_+)r_+)r^3\\
&+r_+(-3r^3(r-r_+)(10\Omega r^4-2(\Omega r_++2)r^3+2r_+(4-3\Omega r_+)r^2+r_+^2(3-2\Omega r_+)r-9r_+^3)\mu_0^6\\
&+\Omega r(12(2\Omega^3r^3-14\Omega r+3)r^5+4(\Omega r(\Omega r(6\Omega r+5)+48)-30)r_+r^4\\
&+2(\Omega r(2\Omega r+21)+50)r_+^2r^3+(83-102\Omega r)r_+^3r^2-3(8\Omega r+59)r_+^4r+48r_+^5)\mu_0^4\\
&+2\Omega^2(12(\Omega r(2\Omega r(3\Omega r+1)-7)-2)r^5+4(\Omega r(\Omega r(16\Omega r+13)+18)+12)r_+r^4\\
&+2(\Omega r(2\Omega r(2\Omega r-1)+13)-14)r_+^2r^3-(176\Omega r+5)r_+^3r^2+3(10\Omega r-27)r_+^4r+24r_+^5)\mu_0^2\\
&+8\Omega^3(12\Omega(2\Omega r(\Omega r+1)+1)r^5+4\Omega(\Omega r(4\Omega r+3)-6)r_+r^4\\
&+2(\Omega r(2\Omega r(2\Omega r-3)-5)-6)r_+^2r^2-(20\Omega r+3)r_+^3r+3(2\Omega r-1)r_+^4))\Bigg)H,
\end{aligned}
\tag{C7}
$$

$$\begin{aligned}
h_2^{vr} \; = \; & \frac{2(r\mu_0^2+2\Omega)(r-r_+)r_+^2}{(r\mu_0^2+4\Omega)(\mu_0^2r^3+r_+)}h_1^{v\chi\prime} + \frac{4r(r\mu_0^2+\Omega(\mu_0^2r^2+2\Omega r+4))r_+^2-2(r\mu_0^2+6\Omega)r_+^3}{r(r\mu_0^2+4\Omega)(\mu_0^2r^3+r_+)}h_1^{v\chi} \\
& +\frac{1}{2}\left(\frac{r(\mu_0^2(2r-3r_+)-4\Omega^2r)}{\Omega(\mu_0^2r^3+r_+)}-2\right)H_2 + \frac{4\Omega^2r^4+r_+(3r_+-4r)}{4(r-r_+)(\mu_0^2r^3+r_+)}H_{2-} \\
& -\frac{1}{24\Omega r^3(r\mu_0^2+4\Omega)(r-r_+)^2(\mu_0^2r^3+r_+)^2} \\
& \times\Biggl(6\mu_2^2\Omega(r\mu_0^2+4\Omega)(r-r_+)(4\Omega^2r^4+r_+(3r_+-4r))r^6 \\
& +r_+\Biggl[r^3(r-r_+)\biggl(24\Omega(\Omega r(\Omega r-1)+1)r^5+4(\Omega r(\Omega r(2\Omega r+3)-15)+6)r_+r^3 \\
& -2(\Omega r(6\Omega r-17)+15)r_+^2r^2-3(2\Omega r+9)r_+^3r+3(9-16\Omega r)r_+^4\biggr)\mu_0^4 \\
& +\Omega r\biggl(24\Omega(\Omega r(2\Omega r(\Omega r+3)-3)+4)r^7+16(\Omega r(\Omega r(2\Omega r(\Omega r-2)+3)-24)+6)r_+r^5 \\
& +2(2\Omega r(4\Omega r(\Omega r(\Omega r-5)-8)+79)-105)r_+^2r^4-4(\Omega r(19\Omega r+16)-42)r_+^3r^3 \\
& +3(4\Omega r(7\Omega r-1)+39)r_+^4r^2+3(16\Omega r-81)r_+^5r+102r_+^6\biggr)\mu_0^2 \\
& +4\Omega^2\biggl(24\Omega^2(2\Omega r(\Omega r+1)+1)r^8+16\Omega(2\Omega^3r^3-6\Omega r-3)r_+r^6 \\
& +2\bigl(2\Omega r(4\Omega r(\Omega r(\Omega r-3)-2)+9)+3\bigr)r_+^2r^4+4(\Omega r(6-7\Omega r)+9)r_+^3r^3 \\
& +3(2\Omega r-1)(6\Omega r+1)r_+^4r^2-51r_+^5r+30r_+^6\biggr)\Biggr]\Biggr)H_- \\
& +\frac{1}{12\Omega^2r^2(r\mu_0^2+4\Omega)(r-r_+)(\mu_0^2r^3+r_+)^2} \\
& \times\Biggl(6\mu_2^2\Omega(r\mu_0^2+4\Omega)(r-r_+)(4\Omega^2r^4+(2r-3r_+)r_+)r^3 \\
& +r_+\Biggl[-3r^3(r-r_+)\biggl(10\Omega r^4-2(\Omega r_++2)r^3 \\
& +2r_+(4-3\Omega r_+)r^2+r_+^2(3-2\Omega r_+)r-9r_+^3\biggr)\mu_0^6 \\
& +\Omega r\biggl(12(2\Omega^3r^3-14\Omega r+3)r^5+4(\Omega r(\Omega r(6\Omega r+5)+48)-30)r_+r^4 \\
& +2(\Omega r(2\Omega r+21)+50)r_+^2r^3+(83-102\Omega r)r_+^3r^2-3(8\Omega r+59)r_+^4r+48r_+^5\biggr)\mu_0^4 \\
& +2\Omega^2\biggl(12(\Omega r(2\Omega r(3\Omega r+1)-7)-2)r^5+4(\Omega r(\Omega r(16\Omega r+13)+18)+12)r_+r^4 \\
& +2(\Omega r(2\Omega r(2\Omega r-1)+13)-14)r_+^2r^3 \\
& -(176\Omega r+5)r_+^3r^2+3(10\Omega r-27)r_+^4r+24r_+^5\biggr)\mu_0^2 \\
& +8\Omega^3\biggl(12\Omega(2\Omega r(\Omega r+1)+1)r^5+4\Omega(\Omega r(4\Omega r+3)-6)r_+r^4 \\
& +2(\Omega r(2\Omega r(2\Omega r-3)-5)-6)r_+^2r^2-(20\Omega r+3)r_+^3r+3(2\Omega r-1)r_+^4\biggr)\Biggr]\Biggr)H, \qquad \text{(C8)}
\end{aligned}$$

$$
\begin{aligned}
h_2^{\theta\theta} \;=\; & \frac{2\Omega r_+(r_+-r)}{3r^2(\mu_0^2 r+4\Omega)} h_1^{v\chi\prime} + \frac{2r_+(\mu_0^2 r r_+ - 2\Omega^2 r^2 + 3\Omega r_+)}{3r^3(\mu_0^2 r+4\Omega)} h_1^{v\chi} + \frac{1}{2r^2} H_{2-} \\
& + \frac{r_+}{6\Omega r^4(r-r_+)(\mu_0^2 r+4\Omega)(\mu_0^2 r^3+r_+)} \\
& \times \Bigg(\mu_0^4 r^2 \Big(6r^3(\Omega r-1) - 2r r_+^2(\Omega r+4) + 13r^2 r_+ + 3r_+^3 \Big) \\
& + 2\mu_0^2\Omega \Big(6r^4(3\Omega r-2) + r^3 r_+(4\Omega r+21) - 2r^2 r_+^2(3\Omega r+2) + 9r r_+^3 - 6r_+^4 \Big) \\
& + 8\Omega^2 \Big(6\Omega r^4 - 2r_+^3(\Omega r+1) + 2r r_+^2(\Omega r+1) + 3r^2 r_+ \Big) \Bigg) H \\
& + \frac{r_+}{12r^5(r-r_+)^2(\mu_0^2 r+4\Omega)(\mu_0^2 r^3+r_+)} \\
& \times \Bigg(-2\mu_0^4 r^4(r-r_+)(r+r_+)(3r^2 - r r_+ - r_+^2) \\
& + \mu_0^2 r \Big(-12\Omega r^6(\Omega r+2) + 2r^4 r_+(8\Omega r+3) + r^3 r_+^2(4\Omega r(\Omega r+6)-15) + 2r^2 r_+^3(9-10\Omega r) \\
& + r r_+^4(4\Omega r-5) - 2r_+^5 \Big) + 4\Omega \Big(-12\Omega^2 r^7 - r^3 r_+^2(2\Omega r(2\Omega r+1)+5) \\
& + 2r^2 r_+^3(\Omega r(2\Omega r+1)+5) + 6r^4 r_+ - 12 r r_+^4 + 4r_+^5 \Big) \Bigg) H_- .
\end{aligned}
\tag{C9}
$$

4. A_H, A_{H_-}, B_H, B_{H_-}, C_H and C_{H_-}

$$
\begin{aligned}
& A_H(r; r_+, \mu_0, \Omega) \\
& := \frac{r_+}{2\Omega r^3(r-r_+)^3\left(\mu_0^2 r+4\Omega\right)\left(\mu_0^2 r^3+r_+\right)} \Big(\mu_0^6 r^4 r_+(r-3r_+)(r-r_+) \\
& + 2\mu_0^4 r\big(-2r^4(\Omega r(\Omega r+1)-3) - r^3 r_+(\Omega r(4\Omega r+1)+6) \\
& + 2r^2 r_+^2(\Omega r-7)(\Omega r+1) + 9r r_+^3(\Omega r+2) - 6r_+^4 \big) \\
& - 8\mu_0^2\Omega\big(r^4(\Omega r+1)(3\Omega r-2) + r^3 r_+(\Omega r(6\Omega r+11)-3) \\
& + r^2 r_+^2(19-\Omega r(3\Omega r+2)) - r r_+^3(\Omega r+14) + 3r_+^4 \big) \\
& + 16\Omega^2\left(-2\Omega r^4(\Omega r+1) - 4\Omega r^3 r_+(\Omega r+2) + r_+^3(3-2\Omega r) + r r_+^2(2\Omega r(\Omega r+3)-5) \right) \Big),
\end{aligned}
\tag{C10}
$$

$$
\begin{aligned}
&A_{H_-}(r;r_+,\mu_0,\Omega) := \\
&\frac{r_+}{4r^4(r-r_+)^4\left(\mu_0^2 r+4\Omega\right)\left(\mu_0^2 r^3+r_+\right)}\Big(\mu_0^4 r^4(r-r_+)\big(8\Omega r^4-2r^3(\Omega r_+ +8) \\
&+2r^2r_+(3\Omega r_+ +13)+17rr_+^2-21r_+^3\big) \\
&+4\mu_0^2 r\big(2\Omega r^6(\Omega r-1)(\Omega r+6)+r^4r_+(\Omega r(\Omega r(4\Omega r+1)+34)-10) \\
&-2r^3r_+^2(\Omega r(\Omega r(\Omega r+3)+8)-6)-r^2r_+^3(\Omega r(\Omega r+19)-10)+2rr_+^4(6\Omega r-11)+9r_+^5\big) \\
&+8\Omega\big(4\Omega^2r^7(\Omega r+1)+2r^4r_+(\Omega r(\Omega r(4\Omega r+7)-1)-5) \\
&-2r^3r_+^2(\Omega r+1)(2\Omega r(\Omega r+3)-3)+3r^2r_+^3(2\Omega r(\Omega r+2)+7) \\
&-rr_+^4(6\Omega r+31)+12r_+^5\big)\Big),
\end{aligned}
\tag{C11}
$$

$$
\begin{aligned}
&B_H(r;r_+,\mu_0,\mu_2,\Omega) := \\
&\frac{1}{6\Omega^2r^3(r\mu_0^2+4\Omega)(r-r_+)(\mu_0^2r^3+r_+)^2}(6\mu_2^2\Omega(r\mu_0^2+4\Omega)(r-r_+)(\mu_0^4r^6+4\Omega^2r^4 \\
&+2\mu_0^2r_+r^3+2(r-r_+)r_+)r^3+r_+(r^6(r-r_+)(6(\Omega r+1)r^2+(2\Omega r-3)r_+r-9r_+^2)\mu_0^8 \\
&+r^3(6(\Omega r(2\Omega r(\Omega r+2)-1)+2)r^4+2(\Omega r+3)(2\Omega r(2\Omega r-1)-5)r_+r^3 \\
&+(\Omega r(4\Omega r(\Omega r+1)+21)+6)r_+^2r^2+(43\Omega r+30)r_+^3r-18(\Omega r+1)r_+^4)\mu_0^6 \\
&+4\Omega r(3(8\Omega^3r^3-14\Omega r+3)r^5+4(\Omega r(\Omega r(5\Omega r+16)+8)-6)r_+r^4 \\
&+(2\Omega r(\Omega r(2\Omega r+11)+30)+9)r_+^2r^3-2(\Omega r-2)(\Omega r+6)r_+^3r^2-6(3\Omega r+5)r_+^4r+9r_+^5)\mu_0^4 \\
&+4\Omega^2r(6(\Omega r(2\Omega r(5\Omega r+1)-7)-2)r^4+4(\Omega r(2\Omega r+3)(7\Omega r+5)+6)r_+r^3 \\
&+(\Omega r(2\Omega r+9)(2\Omega r+11)-36)r_+^2r^2+(42-\Omega r(8\Omega r+87))r_+^3r+12(\Omega r-2)r_+^4)\mu_0^2 \\
&+16\Omega^3(6\Omega(2\Omega r(\Omega r+1)+1)r^5 \\
&+4\Omega(\Omega r(2\Omega r+3)-3)r_+r^4+\Omega(4\Omega^2r^2+15)r_+^2r^3-3(\Omega r-2)r_+^3r-6r_+^4))),
\end{aligned}
\tag{C12}
$$

$$
\begin{aligned}
&B_{H_-}(r;r_+,\mu_0,\mu_2,\Omega) := \\
&\frac{1}{12\Omega r^4(r\mu_0^2+4\Omega)(r-r_+)^2(\mu_0^2r^3+r_+)^2}(r_+(-3\mu_0^6(r-r_+)(8\Omega^2(r+r_+)r^4 \\
&+2\Omega(-3r^3+3r_+r^2+3r_+^2r+r_+^3)r+(4r-3r_+)r_+(r+r_+))r^6 \\
&+\mu_0^4(-24\Omega(\Omega r(\Omega r(\Omega r+5)-4)+1)r^6-4(\Omega r(\Omega r(\Omega r(6\Omega r+13)+54)-18)+6)r_+r^4 \\
&+2(\Omega r(3\Omega r(14\Omega r+5)-31)+21)r_+^2r^3+(\Omega r(2\Omega r(8\Omega r+39)+57)+6)r_+^3r^2 \\
&+(\Omega r(24\Omega r-1)-42)r_+^4r+6(3-4\Omega r)r_+^5)r^3+2\mu_0^2\Omega(-12\Omega(3\Omega r(2\Omega r(\Omega r+1)-1)+4)r^7 \\
&-4(\Omega r(\Omega r(\Omega r(16\Omega r+29)+24)-66)+12)r_+r^5 \\
&-2(\Omega r(\Omega r(2\Omega r(2\Omega r+3)-21)+86)-45)r_+^2r^4+(\Omega r(16\Omega r(2\Omega r+7)-27)-50)r_+^3r^3 \\
&+(\Omega r(1-34\Omega r)-50)r_+^4r^2+2(12\Omega r+47)r_+^5r-42r_+^6)r+8\Omega^2(-12\Omega^2(2\Omega r(\Omega r+1)+1)r^8 \\
&+4\Omega(\Omega r(12-\Omega r(4\Omega r+3))+6)r_+r^6-2(\Omega r(\Omega r(2\Omega r(2\Omega r-3)+9)+12)-3)r_+^2r^4 \\
&+(\Omega r(4\Omega r-9)-2)r_+^3r^3+(5\Omega r(3-2\Omega r)-14)r_+^4r^2+16r_+^5r-6r_+^6)) \\
&-6\mu_2^2\Omega r^6(r\mu_0^2+4\Omega)(r-r_+)(4\Omega^2r^4+r_+(3r_+-4r))),
\end{aligned}
\tag{C13}
$$

$$
\begin{aligned}
&C_H(r;r_+,\mu_0,\mu_2,\Omega) := \\
&\frac{1}{12\Omega r^3(r\mu_0^2+4\Omega)(r-r_+)^2(\mu_0^2r^3+r_+)^2}(6\mu_2^2\Omega(r\mu_0^2+4\Omega)(r-r_+)(4\Omega^2r^4+(2r-3r_+)r_+)r^5 \\
&+r_+(-r^5(r-r_+)(6(7\Omega r-4)r^3-2(\Omega r-15)r_+r^2+(51-10\Omega r)r_+^2r-3(2\Omega r+21)r_+^3)\mu_0^6 \\
&+r^2(12\Omega(2\Omega^3r^3-20\Omega r+7)r^6+4(\Omega r(\Omega r(\Omega r(6\Omega r+5)+56)-54)+3)r_+r^4 \\
&+2(\Omega r(\Omega r(2\Omega r+1)+34)-9)r_+^2r^3+(\Omega r(243-70\Omega r)-36)r_+^3r^2 \\
&+(78-\Omega r(24\Omega r+353))r_+^4r+12(10\Omega r-3)r_+^5)\mu_0^4+2\Omega(12\Omega(\Omega r(2\Omega r(3\Omega r+1)-11)-2)r^7 \\
&+4(\Omega r(\Omega r(\Omega r(16\Omega r+13)+18)-3)+6)r_+r^5 \\
&+2(\Omega r(\Omega r(2\Omega r(2\Omega r-1)+37)+2)-21)r_+^2r^4-(\Omega r(272\Omega r+81)+20)r_+^3r^3 \\
&+(\Omega r(78\Omega r-25)+82)r_+^4r^2+4(6\Omega r-23)r_+^5r+36r_+^6)\mu_0^2+8\Omega^2(12\Omega^2(2\Omega r(\Omega r+1)+1)r^7 \\
&+4\Omega(\Omega r(\Omega r(4\Omega r+3)-6)-3)r_+r^5+2(\Omega r(\Omega r(2\Omega r(2\Omega r-3)-5)-6)-3)r_+^2r^3 \\
&+(\Omega r(9-20\Omega r)+4)r_+^3r^2+(3\Omega r(2\Omega r-9)-14)r_+^4r+2(6\Omega r+5)r_+^5))),
\end{aligned}
\tag{C14}
$$

$$
\begin{aligned}
&C_{H_-}(r;r_+,\mu_0,\mu_2,\Omega) := \\
&\frac{1}{24r^4(r\mu_0^2+4\Omega)(r-r_+)^3(\mu_0^2r^3+r_+)^2}(r_+(-12\mu_0^6(2r-3r_+)(r-r_+)^2(r+r_+)r^8 \\
&-\mu_0^4(r-r_+)(24\Omega(\Omega r(\Omega r-2)+5)r^5+4(\Omega r(\Omega r(2\Omega r-3)-51)+24)r_+r^3 \\
&-2(\Omega r(6\Omega r+7)+48)r_+^2r^2+3(46\Omega r-47)r_+^3r+15(9-8\Omega r)r_+^4)r^5 \\
&+\mu_0^2(-24\Omega^2(\Omega r+4)(2\Omega r(\Omega r-1)+1)r^8-8\Omega(\Omega r(2\Omega r(2\Omega r(\Omega r-2)+3)-51)+48)r_+r^6 \\
&+2(\Omega r(2\Omega r(4\Omega r(\Omega r(5-\Omega r)+2)-79)+381)-24)r_+^2r^4 \\
&+2(2\Omega r(\Omega r(67\Omega r+22)-66)+39)r_+^3r^3-3(3\Omega r(4\Omega r(5\Omega r+1)+85)-20)r_+^4r^2 \\
&-3(\Omega r(16\Omega r-305)+54)r_+^5r+6(12-53\Omega r)r_+^6)r^2+4\Omega(-24\Omega^3(2\Omega r(\Omega r+1)+1)r^{10} \\
&+8\Omega^2(-4\Omega^3r^3+12\Omega r+9)r_+r^8-2(\Omega r(2\Omega r(4\Omega r(\Omega r(\Omega r-3)-2)+9)+3)+24)r_+^2r^5 \\
&+3\Omega(4\Omega r(5-3\Omega r)+5)r_+^4r^4+2(2\Omega r(\Omega r(7\Omega r-12)-9)+39)r_+^3r^4 \\
&-3(\Omega r(8\Omega r-13)+38)r_+^5r^2+6(19-5\Omega r)r_+^6r-36r_+^7)) \\
&-6\mu_2^2\Omega r^8(r\mu_0^2+4\Omega)(r-r_+)(4\Omega^2r^4+r_+(3r_+-4r))).
\end{aligned}
\tag{C15}
$$

5. Expansion of master variables near horizon for $\Omega \neq 0$

$$
\begin{aligned}
H &= \frac{\Omega r_+(2\Omega r_+-1)}{(r-r_+)\left(2\mu_0^2r_++4\Omega\right)}+\frac{\mu_0^2\Omega r_+^2+\Omega}{2\mu_0^2r_++4\Omega} \\
&\quad-\frac{\Omega(r-r_+)\left(-2\Omega r_+\left(\mu_0^4r_+^4-7\right)-3\mu_0^4r_+^4+8\mu_0^2r_+^2+4\Omega^2r_+^2+4\right)}{8r_+(\Omega r_++1)(2\Omega r_++1)\left(\mu_0^2r_++2\Omega\right)}+\mathcal{O}((r-r_+)^2), \qquad \text{(C16)} \\
H_- &= 1+\frac{(r-r_+)\left(\mu_0^4r_+^3-4\mu_0^2r_+-8\Omega^2r_+-6\Omega\right)}{r_+(2\Omega r_++1)\left(\mu_0^2r_++2\Omega\right)} \\
&\quad+\frac{(r-r_+)^2}{4r_+^2(\Omega r_++1)(2\Omega r_++1)\left(\mu_0^2r_++2\Omega\right)}(\Omega^2\left(108r_+-8\mu_0^2r_+^3\right)+\Omega\left(-2\mu_0^4r_+^4+34\mu_0^2r_+^2+48\right) \\
&\quad+\mu_0^2r_+\left(\mu_0^4r_+^4-8\mu_0^2r_+^2+32\right)+48\Omega^3r_+^2)+\mathcal{O}((r-r_+)^2), \qquad \text{(C17)}
\end{aligned}
$$

$$\begin{aligned}
h_1^{v\chi} \;=\;& \beta + \frac{(r-r_+)}{4r_+^2(\Omega r_+ + 1)^2(2\Omega r_+ + 1)(2\Omega r_+ + 3)(\mu_0^2 r_+ + 2\Omega)(\mu_0^2 r_+ + 4\Omega)} \\
& (8\beta\mu_0^6\Omega^3 r_+^8 - 4\Omega r_+(4\Omega(9\beta + 70\Omega) + 109\mu_0^2) - 2r_+^2(2\mu_0^2\Omega(24\beta + 209\Omega) \\
& +8\Omega^3(51\beta + 58\Omega) + 47\mu_0^4) + 8\beta\mu_0^4\Omega^2 r_+^7(3\mu_0^2 + 2\Omega^2) - 2r_+^3(\beta(6\mu_0^4 + 248\mu_0^2\Omega^2 + 816\Omega^4) \\
& +53\mu_0^4\Omega + 210\mu_0^2\Omega^3 + 112\Omega^5) + \mu_0^2 r_+^6(2\mu_0^4\Omega(11\beta + 5\Omega) + 8\mu_0^2\Omega^3(4\beta + \Omega) - 192\beta\Omega^5 - \mu_0^6) \\
& +r_+^4(-2\mu_0^4\Omega(16\beta + 3\Omega) - 24\mu_0^2\Omega^3(38\beta + 3\Omega) + 64\Omega^5(\Omega - 21\beta) + 19\mu_0^6) \\
& +r_+^5(\beta(6\mu_0^6 - 4\mu_0^4\Omega^2 - 704\mu_0^2\Omega^4 - 384\Omega^6) + 19\mu_0^6\Omega + 24\mu_0^4\Omega^3 + 32\mu_0^2\Omega^5) - 408\Omega^2) \\
& +\mathcal{O}((r-r_+)^2),
\end{aligned} \tag{C18}$$

$$\begin{aligned}
H_{2-} \;=\;& \frac{(r-r_+)}{3r_+(\Omega r_+ + 1)(2\Omega r_+ + 1)^2(\mu_0^2 r_+ + 2\Omega)^2(\mu_0^2 r_+ + 4\Omega)} \\
& (\mu_0^6 r_+^3(12\beta r_+ + \mu_0^2 r_+^2 + 3\mu_2^2 r_+^2 - 21) - 8\Omega^5 r_+^2(-24\beta r_+(3\mu_0^2 r_+^2 + 4) + 2\mu_0^4 r_+^4 - 21\mu_0^2 r_+^2 \\
& -24\mu_2^2 r_+^2 - 62) - 4\Omega^4 r_+(-24\beta r_+(3\mu_0^4 r_+^4 + 12\mu_0^2 r_+^2 + 5) + \mu_0^6 r_+^6 + 6\mu_0^4 r_+^4 \\
& -\mu_0^2 r_+^2(36\mu_2^2 r_+^2 + 125) - 60\mu_2^2 r_+^2 - 64) + 2\Omega^3(24\beta r_+(\mu_0^6 r_+^6 + 12\mu_0^4 r_+^4 + 15\mu_0^2 r_+^2 + 2) \\
& -13\mu_0^6 r_+^6 + \mu_0^4(24\mu_2^2 r_+^6 + 22r_+^4) + 6\mu_0^2 r_+^2(17\mu_2^2 r_+^2 + 22) + 12\mu_2^2 r_+^2 - 8) \\
& +\mu_0^2\Omega r_+^2(\mu_0^4 r_+^2(60\beta r_+ + 9\mu_2^2 r_+^2 - 19) + 8\mu_0^2(9\beta r_+ + 3\mu_2^2 r_+^2 - 11) - 6\mu_0^6 r_+^4 - 6\mu_2^2) \\
& +2\Omega^2 r_+(\mu_0^6 r_+^4(48\beta r_+ + 3\mu_2^2 r_+^2 - 25) + 3\mu_0^4 r_+^2(60\beta r_+ + 12\mu_2^2 r_+^2 + 5) \\
& +\mu_0^2(72\beta r_+ + 27\mu_2^2 r_+^2 - 52) - 2\mu_0^8 r_+^6 - 12\mu_2^2) + 32\Omega^6 r_+^3(12\beta r_+ + 7)) \\
& +\mathcal{O}((r-r_+)^2),
\end{aligned} \tag{C19}$$

$$\begin{aligned}
H_2 \;=\;& \frac{\Omega r_+(\Omega(4\mu_0^2 r_+^2 - 6\mu_2^2 r_+^2 + 6) + r_+(4\mu_0^2 + 3\mu_2^2) + 4\Omega^3 r_+^2 + 10\Omega^2 r_+)}{6(r-r_+)(\mu_0^2 r_+ + 2\Omega)^2} \\
& + \frac{1}{6(\Omega r_+ + 1)(\mu_0^2 r_+ + 2\Omega)^2(\mu_0^2 r_+ + 4\Omega)}(6\Omega^4 r_+(8\beta r_+(2\mu_0^2 r_+^2 + 3) - 5\mu_0^2 r_+^2 + 4\mu_2^2 r_+^2 - 40) \\
& +3\mu_0^2\Omega r_+^2(4\beta\mu_0^2 r_+ - \mu_2^2) - 4\Omega^5 r_+^2(-24\beta r_+ + \mu_0^2 r_+^2 + 18) \\
& +2\Omega^3(12\beta r_+(\mu_0^4 r_+^4 + 6\mu_0^2 r_+^2 + 2) + 3\mu_0^2 r_+^2(\mu_2^2 r_+^2 - 40) + 6\mu_2^2 r_+^2 - 56) \\
& +\Omega^2 r_+(12\mu_0^4 r_+^2(3\beta r_+ - 5) + \mu_0^2(48\beta r_+ + 3\mu_2^2 r_+^2 - 82) - 12\mu_2^2) \\
& +6\mu_0^6 r_+^3 - 16\Omega^6 r_+^3) + \mathcal{O}(r-r_+).
\end{aligned} \tag{C20}$$

[1] R. Gregory and R. Laflamme, Phys.Rev.Lett. **70**, 2837 (1993), arXiv:hep-th/9301052, *Black strings and p-branes are unstable.*

[2] R. Gregory and R. Laflamme, Nucl.Phys. **B428**, 399 (1994), arXiv:hep-th/9404071, *The Instability of charged black strings and p-branes.*

[3] O. J. Dias, P. Figueras, R. Monteiro, H. S. Reall, and J. E. Santos, JHEP **1005**, 076 (2010), arXiv:1001.4527, *An instability of higher-dimensional rotating black holes.*

[4] L. Lehner and F. Pretorius, Phys.Rev.Lett. **105**, 101102 (2010), arXiv:1006.5960, *Black Strings, Low Viscosity Fluids, and Violation of Cosmic Censorship.*

[5] E. Thorne, Kip S., E. Price, R.H., and E. Macdonald, D.A., (1986), *BLACK HOLES: THE MEMBRANE PARADIGM.*

[6] S. Bhattacharyya, V. E. Hubeny, S. Minwalla, and M. Rangamani, JHEP **0802**, 045 (2008), arXiv:0712.2456, *Nonlinear Fluid Dynamics from Gravity.*

[7] R. Emparan, T. Harmark, V. Niarchos, and N. A. Obers, Phys.Rev.Lett. **102**, 191301 (2009), arXiv:0902.0427, *World-Volume Effective Theory for Higher-Dimensional Black Holes.*

[8] R. Emparan, T. Harmark, V. Niarchos, and N. A. Obers, JHEP **1003**, 063 (2010), arXiv:0910.1601, *Essentials of Blackfold Dynamics.*

[9] V. Cardoso and O. J. Dias, Phys.Rev.Lett. **96**, 181601 (2006), arXiv:hep-th/0602017, *Rayleigh-Plateau and Gregory-Laflamme instabilities of black strings.*

[10] V. Cardoso and L. Gualtieri, Class.Quant.Grav. **23**, 7151 (2006), arXiv:hep-th/0610004, *Equilibrium configurations of fluids and their stability in higher dimensions.*

[11] J. Camps, R. Emparan, and N. Haddad, JHEP **1005**, 042 (2010), arXiv:1003.3636, *Black Brane Viscosity and the Gregory-Laflamme Instability.*

[12] B. Kleihaus, J. Kunz, and E. Radu, JHEP **0705**, 058 (2007), arXiv:hep-th/0702053, *Rotating nonuniform black string solutions.*

[13] R. Emparan and R. C. Myers, JHEP **0309**, 025 (2003), arXiv:hep-th/0308056, *Instability of ultra-spinning black holes.*

[14] M. M. Caldarelli, O. J. Dias, R. Emparan, and D. Klemm, JHEP **0904**, 024 (2009), arXiv:0811.2381, *Black Holes as Lumps of Fluid.*

[15] A. M. Karnik, Astrophysics and Space Science **83**, 209 (1982), *Effect of rotation on the stability of a gravitating cylinder.*

[16] T. Regge and J. A. Wheeler, Phys.Rev. **108**, 1063 (1957), *Stability of a Schwarzschild singularity.*

[17] F. J. Zerilli, Journal of Mathematical Physics **11**, 2203 (1970), *Tensor Harmonics in Canonical Form for Gravitational Radiation and Other Applications.*

[18] F. Zerilli, Phys.Rev. **D2**, 2141 (1970), *Gravitational field of a particle falling in a schwarzschild geometry analyzed in tensor harmonics.*